

Dynamically generated resonances

E. OSET¹, R. MOLINA¹, D. NICMORUS², L.S. GENG³, P. GONZALEZ¹, J. VIJANDE⁴, S. SARKAR⁵, BAO XI SUN⁶, M. J. VICENTE VACAS¹, A. RAMOS⁷, E. J. GARZON¹, A. MARTINEZ TORRES⁸, K. KHEMCHANDANI⁹

¹*Departamento de Física Teórica and IFIC, Universidad de Valencia-CSIC,
E-46071 Valencia, Spain*

²*Fachbereich Theoretische Physik, Institut für Physik, Karl-Franzens-Universitaet
Graz, Universitaetsplatz 5, A-8010 Graz, Austria*

³*Physik-Department, Technische Universitaet Muenchen, D-85747 Garching,
Germany*

⁴*Departamento de Física Atomica Molecular y Nuclear and IFIC, Centro Mixto
Universidad de Valencia-CSIC*

⁵*Variable Energy Cyclotron Centre, 1/AF, Bidhannagar, Kolkata 700064, India*

⁶*Institute of Theoretical Physics, College of Applied Sciences, Beijing University of
Technology, Beijing 100124, China*

⁷*Departament d'Estructura i Constituents de la Matèria and Institut de Ciències
del Cosmos, Universitat de Barcelona, 08028 Barcelona, Spain*

⁸*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502,
Japan*

⁹*Research Center for Nuclear Physics (RCNP), Osaka University, Ibaraki, Osaka
567-0047, Japan*

We present recent results on the vector meson-vector meson and vector meson-baryon interaction using a unitary approach based on the hidden-gauge Lagrangians. For the vector-vector case we find that 11 states get dynamically generated, corresponding to poles of the scattering matrices on the second Riemann sheet. In the vector-baryon sector we also find 9 states dynamically generated from the vector-baryon octet interaction and 10 from the vector-baryon decuplet interaction. We also report on baryon states found from the interaction of two mesons and a baryon.

§1. Introduction

Early in 1978, S. Weinberg made a great gift to hadron physics by showing that the dynamics of QCD in the hadron world can be addressed at low energies by means of effective theories in which the building blocks are the ground state mesons and baryons.¹⁾ This idea has produced tools to address the interaction of mesons or mesons and baryons, mainly through chiral Lagrangians, which have had a tremendous impact on our understanding of the spectrum of meson and baryon resonances. We all accept that ground states of mesons and baryons are made of $q\bar{q}$ or three q respectively. Yet, the spectrum of excited hadronic states can be much richer.

The building blocks in these chiral theories are the low energy hadrons, such as the proton and baryons of its SU(3) octet. To these one adds also the decuplet of

the Δ , considered as spin realignments of the three quark ground state. The basic mesons are the pion and mesons of its octet, to which one also adds the nonet of the ρ , which also corresponds to spin realignments of the $q\bar{q}$ ground state.

What about meson and baryon resonances? The logical answer is that they are excitations of the quarks, which is the essence of quark models. Yet, although extensive theoretical efforts are spent by QCD motivated approaches which describe hadron resonances as excitations of quarks,^{2),3)} a complete and reliable calculation of the entire excited hadron spectrum is still missing.

It is interesting to recall basic facts from the baryon spectrum. The first excited N^* states are the $N^*(1440)$ ($1/2^+$) and the $N^*(1535)$ ($1/2^-$). In quark models these states require a quark excitation of about 500-600 MeV. If this is the case, one may think that it takes less energy to create one pion, or two (140-280 MeV). The question is whether they can be bound or get trapped in a resonant state. How do we know if this can occur? We need dynamics, a potential for the interaction of mesons with ground state baryons and then solve the Schroedinger equation in coupled channels (or the Bethe Salpeter equation where the mesons are treated relativistically). This information can be extracted from chiral Lagrangians: the effective theory of QCD at low energies. This is the philosophy behind the idea of dynamically generated baryons: Many resonances are generated in this way, like the $1/2^-$ states from meson baryon: $N^*(1535)$,⁴⁾ two $\Lambda(1405)$ ⁵⁾ or the $1/2^+$ states from two mesons and a baryon, like the $N^*(1710)$ and others.^{6),7)}

Similarly, the interaction of pseudoscalar mesons leads to a good description of the low lying scalar mesons $f_0(600)$, $f_0(980)$, and $a_0(980)$ ⁸⁾⁻¹¹⁾

Inspired by the success of the unitary chiral approach, a further extension has recently been taken to study the interaction between two vector mesons and between one vector meson and one baryon.¹²⁾⁻¹⁷⁾ The novelty is that instead of using interaction kernels provided by ChPT, one uses transition amplitudes provided by the hidden-gauge Lagrangians, which lead to a suitable description of the interaction of vector mesons among themselves and of vector mesons with other mesons or baryons. Coupled-channel unitarity works in the same way as in the unitary chiral approach, but now the dynamics is provided by the hidden-gauge Lagrangians.¹⁸⁾⁻²¹⁾ As shown by several recent works,²²⁾⁻²⁴⁾ this combination seems to work very well.

In this talk, we give an overview of recent developments concerning the interaction of vector mesons among themselves and vector mesons with baryons using the unitary approach, where several resonances appear as dynamically generated states.

§2. Theoretical framework

We follow the formalism of the hidden gauge interaction for vector mesons of¹⁸⁾⁻²¹⁾ (see also²⁵⁾ for a practical set of Feynman rules). The Lagrangian involving the interaction of vector mesons amongst themselves is given by

$$\mathcal{L}_{III} = -\frac{1}{4}\langle V_{\mu\nu} V^{\mu\nu} \rangle, \quad (2.1)$$

where the symbol $\langle \rangle$ stands for the trace in the $SU(3)$ space and $V_{\mu\nu}$ is given by

$$V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu - ig[V_\mu, V_\nu] , \quad (2.2)$$

where g is $g = \frac{M_V}{2f}$, with $f = 93$ MeV the pion decay constant. The magnitude V_μ is the ordinary $SU(3)$ matrix of the vectors of the octet of the ρ

The lagrangian \mathcal{L}_{III} gives rise to a contact term coming from $[V_\mu, V_\nu][V_\mu, V_\nu]$, as well as to a three vector vertex

$$\mathcal{L}_{III}^{(c)} = \frac{g^2}{2} \langle V_\mu V_\nu V^\mu V^\nu - V_\nu V_\mu V^\mu V^\nu \rangle ; \quad \mathcal{L}_{III}^{(3V)} = ig \langle (V^\mu \partial_\nu V_\mu - \partial_\nu V_\mu V^\mu) V^\nu \rangle, \quad (2.3)$$

In this case one finds an analogy to the coupling of vectors to pseudoscalars given in the same theory by

$$\mathcal{L}_{VPP} = -ig \langle [P, \partial_\nu P] V^\nu \rangle , \quad (2.4)$$

where P is the $SU(3)$ matrix of the pseudoscalar fields.

In a similar way, one obtains the Lagrangian for the coupling of vector mesons to the baryon octet given by^{26),27) *}

$$\mathcal{L}_{BBV} = g \left(\langle \bar{B} \gamma_\mu [V^\mu, B] \rangle + \langle \bar{B} \gamma_\mu B \rangle \langle V^\mu \rangle \right) \quad (2.5)$$

where B is now the ordinary $SU(3)$ matrix of the baryon octet

With these ingredients we can construct the Feynman diagrams that lead to the $PB \rightarrow PB$ and $VB \rightarrow VB$ interaction, by exchanging a vector meson between the pseudoscalar or the vector meson and the baryon, as depicted in Fig. 1.




Fig. 1. Diagrams contributing to the pseudoscalar-baryon (a) or vector- baryon (b) interaction via the exchange of a vector meson.

From the diagram of Fig. 1(a), and under the low energy approximation of neglecting q^2/M_V^2 in the propagator of the exchanged vector, where q is the momentum transfer, one obtains the same amplitudes as obtained from the ordinary chiral Lagrangian for pseudoscalar-baryon octet interaction, namely the Weinberg-Tomozawa terms. The approximation of neglecting the three momenta of the vectors implies that V^ν in eq. (2.3) corresponds to the exchanged vector and the analogy with eq. (2.4) is more apparent. Note that $\epsilon_\mu \epsilon^\mu$ becomes $-\vec{\epsilon} \vec{\epsilon}'$ and the signs of the Lagrangians also agree.

*) Correcting a misprint in²⁶⁾




Fig. 2. Transition amplitudes V appearing in the coupled-channel Bethe-Salpeter equation.

One can see that the cases with $(I, S) = (3/2, 0)$, $(2, -1)$ and $(3/2, -2)$, the last two corresponding to exotic channels, have a repulsive interaction and do not produce poles in the scattering matrices. However, the sectors $(I, S) = (1/2, 0)$, $(0, -1)$, $(1, -1)$ and $(1/2, -2)$ are attractive and one finds bound states and resonances in these cases.

The scattering matrix is obtained solving the coupled channels Bethe Salpeter equation in the on shell factorization approach of^(28),29)

$$T = [1 - V G]^{-1} V \quad (2.6)$$

with G being the loop function of a vector meson and a baryon. This function is convoluted with the spectral function of the vector mesons to take into account their width as done in.¹⁴⁾

In this case the factor $\vec{\epsilon}\vec{\epsilon}'$, appearing in the potential V , factorizes also in the T matrix for the external vector mesons. This trivial spin structure is responsible for having degenerate states with spin-parity $1/2^-, 3/2^-$ for the interaction of vectors with the octet of baryons and $1/2^-, 3/2^-, 5/2^-$ for the interaction of vectors with the decuplet of baryons.

What we have done here for the interaction of vectors with the octet of baryons can be done for the interaction of vectors with the decuplet of baryons, and the interaction is obtained directly from that of the pseudoscalar-decuplet of baryons studied in.^{30),31)} The study of this interaction in⁽¹⁵⁾⁻¹⁷⁾ leads also to the generation of many resonances which are described below.

We search for poles in the scattering matrices in the second Riemann sheet, as defined in previous works.³²⁾

For the case of vector-vector interaction the procedure followed is similar. We outline the main ingredients of the unitary approach (details can be found in Refs.^{12),13)}). There are two basic building-blocks in this approach: transition amplitudes provided by the hidden-gauge Lagrangians and a unitarization procedure. In Refs.^{12),13)} three mechanisms, as shown in Fig. 2, have been taken into account for the calculation of the transition amplitudes V : the four-vector-contact term, the $t(u)$ -channel vector-exchange amplitude, and the direct box amplitude with two intermediate pseudoscalar mesons.

Among the three mechanisms considered for V , the four-vector-contact term and $t(u)$ -channel vector-exchange one are responsible for the formation of resonances or

bound states if the interaction generated by them is strong enough. In this sense, the dynamically generated states can be thought of as “vector meson-vector meson molecules.” On the other hand, the consideration of the imaginary part of the direct box amplitude allows the generated states to decay into two pseudoscalars. It should be stressed that in the present approach these two mechanisms play quite different roles: the four-vector-contact term and the $t(u)$ -channel vector-exchange one are responsible for generating the resonances whereas the direct box amplitude mainly contributes to their decays.

To calculate the box diagram, one has to further introduce two parameters, Λ and Λ_b . The parameter Λ regulates the four-point loop function, and Λ_b is related to the form factors used for the vector-pseudoscalar-pseudoscalar vertex, which is inspired by the empirical form factors used in the study of vector-meson decays.³³⁾ The values of Λ and Λ_b have been fixed in Ref.¹²⁾ to obtain the widths of the $f_0(1370)$ and $f_2(1270)$. They are found to provide also a good description of the widths of the $f'_2(1525)$, $K_2^*(1430)$, and $f_0(1710)$.

§3. Resonances from the interaction of two vector mesons

Table I. The properties, (mass, width) [in units of MeV], of the 11 dynamically generated states and, if existing, of those of their PDG counterparts.³⁴⁾ The association of the dynamically generated states with their experimental counterparts is determined by matching their mass, width, and decay pattern.

$I^G(J^{PC})$	Theory			PDG data		
	Pole position	Real axis		Name	Mass	Width
		$\Lambda_b = 1.4 \text{ GeV}$	$\Lambda_b = 1.5 \text{ GeV}$			
$0^+(0^{++})$	(1512,51)	(1523,257)	(1517,396)	$f_0(1370)$	1200~1500	200~500
$0^+(0^{++})$	(1726,28)	(1721,133)	(1717,151)	$f_0(1710)$	1724 ± 7	137 ± 8
$0^-(1^{+-})$	(1802,78)	(1802,49)		h_1		
$0^+(2^{++})$	(1275,2)	(1276,97)	(1275,111)	$f_2(1270)$	1275.1 ± 1.2	$185.0^{+2.9}_{-2.4}$
$0^+(2^{++})$	(1525,6)	(1525,45)	(1525,51)	$f'_2(1525)$	1525 ± 5	73^{+6}_{-5}
$1^-(0^{++})$	(1780,133)	(1777,148)	(1777,172)	a_0		
$1^+(1^{+-})$	(1679,235)	(1703,188)		b_1		
$1^-(2^{++})$	(1569,32)	(1567,47)	(1566,51)	$a_2(1700)??$		
$1/2(0^+)$	(1643,47)	(1639,139)	(1637,162)	K_0^*		
$1/2(1^+)$	(1737,165)	(1743,126)		$K_1(1650)?$		
$1/2(2^+)$	(1431,1)	(1431,56)	(1431,63)	$K_2^*(1430)$	1429 ± 1.4	104 ± 4

Searching for poles of the scattering matrix T on the second Riemann sheet, we find 11 states in nine strangeness-isospin-spin channels as shown in Table I. Theoretical masses and widths are obtained with two different methods: In the “pole position” method, the mass corresponds to the real part of the pole position

on the complex plane and the width corresponds to twice its imaginary part. In this case, the box diagrams corresponding to decays into two pseudoscalars are not included. In the "real axis" method, the resonance parameters are obtained from the modulus squared of the amplitudes of the dominant channel of each state on the real axis^{*)}, where the mass corresponds to the energy at which the modulus squared has a maximum and the width corresponds to the difference between the two energies where the modulus squared is half of the maximum value. In this latter case, the box amplitudes are included. The results shown in Table I have been obtained using two different values of Λ_b , which serve to quantify the uncertainties related to this parameter.

Our treatment of the box amplitudes enables us to obtain the decay branching ratios of the generated states into two pseudoscalar mesons using the real-axis method and this is explained in detail in Ref.¹³⁾

It is interesting to note that out of the 21 combinations of strangeness, isospin and spin, we have found resonances only in nine of them. In all the "exotic" channels, from the point of view that they cannot be formed from $q\bar{q}$ combinations, we do not find dynamically generated resonances.

Applications of the results exposed above have been done to study different processes. As an example, the radiative decay into $\gamma\gamma$ of the $f_2(1270)$ and $f_0(1370)$ has been studied in³⁵⁾ and good agreement with experiment is found. Similarly, the decay into $\gamma\gamma$ or a γ and a vector meson for other resonances of¹³⁾ has also been evaluated in,²²⁾ with also good agreement with experiment when available. Another test successfully passed by the former approach to these resonances is the decay of J/ψ into $\phi(\omega)$ and one of the resonances generated in,¹³⁾ as explained in.²³⁾ Another test is the J/ψ decay into a γ and one of the resonances of,¹³⁾ which has been studied in.²⁴⁾

§4. Resonances from the interaction of vector mesons with baryons

In table II we show a summary of the results obtained from the interaction of vectors with the octet of baryons¹⁷⁾ and the tentative association to known states.³⁴⁾

For the $(I, S) = (1/2, 0)$ N^* states there is the $N^*(1700)$ with $J^P = 3/2^-$, which could correspond to the state we find with the same quantum numbers around the same energy. We also find in the PDG the $N^*(1650)$, which could be the near degenerate spin partner of the $N^*(1700)$ that we predict in the theory. It is interesting to recall that in the study of Ref.³⁶⁾ a pole is found around 1700 MeV, with the largest coupling to ρN states. Around 2000 MeV, where we find another N^* resonance, there are the states $N^*(2080)$ and $N^*(2090)$, with $J^P = 3/2^-$ and $J^P = 1/2^-$ respectively, showing a good approximate spin degeneracy.

For the case $(I, S) = (0, -1)$ there is in the PDG one state, the $\Lambda(1800)$ with $J^P = 1/2^-$, remarkably close to the energy where we find a Λ state. The state obtained around 1900 MeV could correspond to the $\Lambda(2000)$ cataloged in the PDG with unknown spin and parity.

^{*)} See Tables I, II, and III of Ref.¹³⁾

Table II. The properties of the 9 dynamically generated resonances and their possible PDG counterparts.

I, S	Theory		PDG data				
	pole position	real axis mass width	name	J^P	status	mass	width
$1/2, 0$	—	1696 92	$N(1650)$	$1/2^-$	***	1645-1670	145-185
			$N(1700)$	$3/2^-$	**	1650-1750	50-150
	$1977 + i53$	1972 64	$N(2080)$	$3/2^-$	**	≈ 2080	180-450
			$N(2090)$	$1/2^-$	*	≈ 2090	100-400
$0, -1$	$1784 + i4$	1783 9	$\Lambda(1690)$	$3/2^-$	***	1685-1695	50-70
			$\Lambda(1800)$	$1/2^-$	**	1720-1850	200-400
	$1907 + i70$	1900 54	$\Lambda(2000)$	$?^?$	*	≈ 2000	73-240
		$2158 + i13$	2158 23				
$1, -1$	—	1830 42	$\Sigma(1750)$	$1/2^-$	**	1730-1800	60-160
	—	1987 240	$\Sigma(1940)$	$3/2^-$	**	1900-1950	150-300
			$\Sigma(2000)$	$1/2^-$	*	≈ 2000	100-450
$1/2, -2$	$2039 + i67$	2039 64	$\Xi(1950)$	$?^?$	**	1950 ± 15	60 ± 20
	$2083 + i31$	2077 29	$\Xi(2120)$	$?^?$	*	≈ 2120	25

The case of the Σ states having $(I, S) = (1, -1)$ is rather interesting. The state that we find around 1830 MeV, could be associated to the $\Sigma(1750)$ with $J^P = 1/2^-$. More interesting seems to be the case of the state obtained around 1990 MeV that could be related to two PDG candidates, again nearly degenerate, the $\Sigma(1940)$ and the $\Sigma(2000)$, with spin and parity $J^P = 3/2^-$ and $J^P = 1/2^-$ respectively.

Finally, for the case of the cascade resonances, $(I, S) = (1/2, -2)$, we find two states, one around 2040 MeV and the other one around 2080 MeV. There are two cascade states in the PDG around this energy region with spin parity unknown, the $\Xi(1950)$ and the $\Xi(2120)$. Although the experimental knowledge of this sector is relatively poor, a program is presently running at Jefferson Lab to improve on this situation.³⁷⁾

The case of the vector interaction with the decuplet is similar³¹⁾ and we show the results in Table III

We also can see that in many cases the experiment shows the near degeneracy predicted by the theory. Particularly, the case of the three Δ resonances around 1920 MeV is very interesting. One observes a near degeneracy in the three spins $1/2^-, 3/2^-, 5/2^-$, as the theory predicts. It is also very instructive to recall that the case of the $\Delta(5/2^-)$ is highly problematic in quark models since it has a $3 \hbar\omega$ excitation and comes out always with a very high mass.^{15),38)}

The association of states found to some resonances reported in the PDG for the case of Λ , Σ and Ξ states looks also equally appealing as one can see from the table.

Table III. The properties of the 10 dynamically generated resonances and their possible PDG counterparts. We also include the N^* bump around 2270 MeV and the Δ^* bump around 2200 MeV.

S, I	Theory			PDG data				
	pole position	real axis		name	J^P	status	mass	width
		mass	width					
0, 1/2	$1850 + i5$	1850	11	$N(2090)$	$1/2^-$	\star	1880-2180	95-414
				$N(2080)$	$3/2^-$	$\star\star$	1804-2081	180-450
		2270(<i>bump</i>)		$N(2200)$	$5/2^-$	$\star\star$	1900-2228	130-400
0, 3/2	$1972 + i49$	1971	52	$\Delta(1900)$	$1/2^-$	$\star\star$	1850-1950	140-240
				$\Delta(1940)$	$3/2^-$	\star	1940-2057	198-460
				$\Delta(1930)$	$5/2^-$	$\star\star\star$	1900-2020	220-500
		2200(<i>bump</i>)		$\Delta(2150)$	$1/2^-$	\star	2050-2200	120-200
-1, 0	$2052 + i10$	2050	19	$\Lambda(2000)$	$?^?$	\star	1935-2030	73-180
-1, 1	$1987 + i1$	1985	10	$\Sigma(1940)$	$3/2^-$	$\star\star\star$	1900-1950	150-300
	$2145 + i58$	2144	57	$\Sigma(2000)$	$1/2^-$	\star	1944-2004	116-413
	$2383 + i73$	2370	99	$\Sigma(2250)$	$?^?$	$\star\star\star$	2210-2280	60-150
				$\Sigma(2455)$	$?^?$	$\star\star$	2455 ± 10	100-140
-2, 1/2	$2214 + i4$	2215	9	$\Xi(2250)$	$?^?$	$\star\star$	2189-2295	30-130
	$2305 + i66$	2308	66	$\Xi(2370)$	$?^?$	$\star\star$	2356-2392	75-80
	$2522 + i38$	2512	60	$\Xi(2500)$	$?^?$	\star	2430-2505	59-150
-3, 0	$2449 + i7$	2445	13	$\Omega(2470)$	$?^?$	$\star\star$	2474 ± 12	72 ± 33

The reasonable results reported here should give a stimulus to search experimentally for the missing spin partners of the already observed states, as well as possible new ones.

One interesting application of these results is the evaluation of the radiative decay of these resonances into a γ and a baryon of the octet or the decuplet, according to the case studied above. Predictions are made in³⁹⁾ for radiative decay widths into γ and a baryon, and, in particular, for the case of decay into γ and a baryon of the octet, the helicity amplitudes are evaluated and compared to data when available. The agreement with experiment is fair, account taken of the experimental uncertainties. A comparison with predictions of quark models is also made in.³⁹⁾

§5. States of two mesons and a baryon

In^{6), 7)} a formalism was developed to study Faddeev equations of systems of two mesons and a stable baryon. The interaction of the pairs was obtained from the chiral

unitary approach, which proves quite successful to give the scattering amplitudes of meson-meson and meson-baryon systems in the region of energies of interest to us. The spectacular finding is that, leaving apart the Roper resonance, whose structure is far more elaborate than originally thought,^{40),41)} all the low lying $J^P = 1/2^+$ excited states are obtained as bound states or resonances of two mesons and one baryon in coupled channels.

Particularly relevant is the issue of a possible bound state of $K\bar{K}N$. In,⁴²⁾ using variational methods, the authors found a bound state of $K\bar{K}N$, with the $K\bar{K}$ being in the $a_0(980)$ state.⁴²⁾ The system was studied a posteriori in⁴³⁾ and it was found to appear at the same energy and the same configuration, although with a mixture of $f_0(980)N$, see fig. 3. This state appears around 1920 MeV with $J^P = 1/2^+$. In a recent paper⁴⁴⁾ some arguments were given to associate this state with the bump that one sees in the $\gamma p \rightarrow K^+ \Lambda$ reaction around this energy, which is clearly visible in recent accurate experiments.^{45),46)} If this association was correct there would be other experimental consequences, as an enhanced strength of the $\gamma p \rightarrow K^+ K^- p$ cross section close to threshold, as well as a shift of strength close to the $K\bar{K}$ threshold in the invariant mass distribution of the kaon pair. This reaction is right now under study.⁴⁷⁾ Another suggestion of⁴⁴⁾ is to measure the total γp spin $S_z = 1/2$ and $S_z = 3/2$ amplitudes, the z -direction along the photon momentum, since this would discriminate the cases where the peak around 1920 MeV is due to a $1/2^+$ or a $3/2^+$ resonance.




Fig. 3. $|T|^2$ for the $NK\bar{K} \rightarrow NK\bar{K}$ transition, showing a peak that is associated to a tentative $N^*(1910)$ resonance.

§6. Acknowledgments

L. S. Geng acknowledges support from the MICINN in the Program Juan de la Cierva. This work is partly supported by DGICYT Contract No. FIS2006-03438, the Generalitat Valenciana in the program PROMETEO and the EU Integrated Infrastructure Initiative Hadron Physics Project under contract RII3-CT-2004-506078.

References

- 1) S. Weinberg, *Physica A* **96**, 327 (1979).
- 2) A. Holl, A. Krassnigg, P. Maris, C. D. Roberts and S. V. Wright, *Phys. Rev. C* **71**, 065204 (2005) [arXiv:nucl-th/0503043].
- 3) A. Krassnigg, *Phys. Rev. D* **80**, 114010 (2009) [arXiv:0909.4016 [hep-ph]].
- 4) N. Kaiser, P. B. Siegel and W. Weise, *Nucl. Phys. A* **594**, 325 (1995)
- 5) D. Jido, J. A. Oller, E. Oset, A. Ramos and U. G. Meissner, *Nucl. Phys. A* **725**, 181 (2003)
- 6) A. Martinez Torres, K. P. Khemchandani and E. Oset, *Phys. Rev. C* **77**, 042203 (2008) .
- 7) K. P. Khemchandani, A. Martinez Torres and E. Oset, *Eur. Phys. J. A* **37**, 233 (2008)
- 8) J. A. Oller, E. Oset and J. R. Pelaez, *Phys. Rev. D* **59**, 074001 (1999) [Erratum-ibid. *D* **60**, 099906 (1999 ERRATUM, D75, 099903, 2007)]
- 9) N. Kaiser, *Eur. Phys. J. A* **3**, 307 (1998).
- 10) J. R. Pelaez and G. Rios, *Phys. Rev. Lett.* **97**, 242002 (2006)
- 11) J. Nieves and E. Ruiz Arriola, *Nucl. Phys. A* **679**, 57 (2000)
- 12) R. Molina, D. Nicmorus and E. Oset, *Phys. Rev. D* **78**, 114018 (2008)
- 13) L. S. Geng and E. Oset, *Phys. Rev. D* **79**, 074009 (2009).
- 14) R. Molina, H. Nagahiro, A. Hosaka and E. Oset, *Phys. Rev. D* **80**, 014025 (2009)
- 15) P. Gonzalez, E. Oset and J. Vijande, *Phys. Rev. C* **79**, 025209 (2009).
- 16) S. Sarkar, B. X. Sun, E. Oset and M. J. V. Vacas, *Eur. Phys. J. A* **44**, 431 (2010).
- 17) E. Oset and A. Ramos, *Eur. Phys. J. A* **44**, 445 (2010) [arXiv:0905.0973 [hep-ph]].
- 18) M. Bando *et al.* *Phys. Rev. Lett.* **54**, 1215 (1985).
- 19) M. Bando, T. Kugo and K. Yamawaki, *Phys. Rept.* **164**, 217 (1988).
- 20) M. Harada and K. Yamawaki, *Phys. Rept.* **381**, 1 (2003)
- 21) U. G. Meissner, *Phys. Rept.* **161**, 213 (1988).
- 22) T. Branz, L. S. Geng and E. Oset, *Phys. Rev. D* **81**, 054037 (2010)
- 23) A. Martinez Torres, L. S. Geng, L. R. Dai, B. X. Sun, E. Oset and B. S. Zou, *Phys. Lett. B* **680**, 310 (2009)
- 24) L. S. Geng, F. K. Guo, C. Hanhart, R. Molina, E. Oset and B. S. Zou, *Eur. Phys. J. A* **44**, 305 (2010)
- 25) H. Nagahiro, L. Roca, A. Hosaka and E. Oset, *Phys. Rev. D* **79**, 014015 (2009) .
- 26) F. Klingl, N. Kaiser and W. Weise, *Nucl. Phys. A* **624** (1997) 527 .
- 27) J. E. Palomar and E. Oset, *Nucl. Phys. A* **716**, 169 (2003) .
- 28) E. Oset and A. Ramos, *Nucl. Phys. A* **635**, 99 (1998)
- 29) J. A. Oller and U. G. Meissner, *Phys. Lett. B* **500**, 263 (2001)
- 30) E. E. Kolomeitsev and M. F. M. Lutz, *Phys. Lett. B* **585** (2004) 243 .
- 31) S. Sarkar, E. Oset and M. J. Vicente Vacas, *Nucl. Phys. A* **750**, 294 (2005) [Erratum-ibid. *A* **780**, 78 (2006)] .
- 32) L. Roca, E. Oset and J. Singh, *Phys. Rev. D* **72**, 014002 (2005) .
- 33) A. I. Titov, B. Kampf and B. L. Reznik, *Eur. Phys. J. A* **7**, 543 (2000); A. I. Titov, B. Kampf and B. L. Reznik, *Phys. Rev. C* **65**, 065202 (2002).
- 34) C. Amsler *et al.* [Particle Data Group], *Phys. Lett. B* **667**, 1 (2008).
- 35) H. Nagahiro, J. Yamagata-Sekihara, E. Oset, S. Hirenzaki and R. Molina, *Phys. Rev. D* **79**, 114023 (2009)
- 36) M. Doring, C. Hanhart, F. Huang, S. Krewald and U. G. Meissner, *Nucl. Phys. A* **829**, 170 (2009)
- 37) B. M. K. Nefkens, *AIP Conf. Proc.* **870**, 405 (2006).
- 38) P. Gonzalez, J. Vijande and A. Valcarce, *Phys. Rev. C* **77**, 065213 (2008) .
- 39) B. X. Sun, E. J. Garzon and E. Oset, *Phys. Rev. D*, in print. arXiv:1003.4664 [hep-ph].
- 40) O. Krehl, C. Hanhart, S. Krewald and J. Speth, *Phys. Rev. C* **62**, 025207 (2000) .
- 41) M. Dillig and M. Schott, *Phys. Rev. C* **75**, 067001 (2007) [Erratum-ibid. *C* **76**, 019903 (2007)] .
- 42) Y. Kanada-En'yo and D. Jido, *Phys. Rev. C* **78**, 025212 (2008) .
- 43) A. Martinez Torres, K. P. Khemchandani and E. Oset, *Phys. Rev. C* **79**, 065207 (2009)
- 44) A. Martinez Torres, K. P. Khemchandani, U. G. Meissner and E. Oset, *Eur. Phys. J. A* **41**, 361 (2009)
- 45) R. Bradford *et al.* [CLAS Collaboration], *Phys. Rev. C* **73**, 035202 (2006) .
- 46) M. Sumihama *et al.* [LEPS Collaboration], *Phys. Rev. C* **73**, 035214 (2006) .
- 47) T. Nakano, talk at the NSTAR2009 Workshop, Beijing, April 2009.